



Left atrial thrombus caused by the primary antiphospholipid syndrome causing critical functional mitral stenosis

S Yusuf, B P Madden and C W Pumphrey

Heart 2003;89;262-
doi:10.1136/heart.89.3.262

Updated information and services can be found at:
<http://heart.bmjournals.com/cgi/content/full/89/3/262>

These include:

Rapid responses

You can respond to this article at:
<http://heart.bmjournals.com/cgi/eletter-submit/89/3/262>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Other imaging techniques](#) (1117 articles)
[Valve diseases](#) (278 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Heart* go to:
<http://www.bmjournals.com/subscriptions/>

CARDIOVASCULAR MEDICINE

Blood culture negative endocarditis: analysis of 63 cases presenting over 25 years

C C Lamas, S J Eykyn

Heart 2003;89:258-262

Objective: To analyse cases of blood culture negative endocarditis (BCNE) seen at St Thomas' Hospital, London, between 1975 and 2000.

Methods: Data on all episodes of endocarditis with negative blood cultures seen at St Thomas' Hospital between 1975 and 2000 were collected prospectively and analysed retrospectively.

Results: Sixty three patients with BCNE were seen during the study period: 48 (76%) with native and 15 (24%) prosthetic valve infection. BCNE accounted for 12.2% of the 516 cases of endocarditis seen at St Thomas' Hospital. The diagnosis of endocarditis was clinically definite by the Duke criteria in only 21% (7 of 34) of cases of pathologically proven native valve endocarditis but in 62% (21 of 34) of cases by the St Thomas' modifications of the criteria. Comparable figures for the 11 cases of pathologically proven prosthetic valve endocarditis were 45% and 73%. Despite negative blood cultures a causative organism was identified in 31 (49%) of the 63 cases: in 15 by serology (8 *Coxiella burnetii*, 6 *Bartonella* species, and 1 *Chlamydia psittaci*); in 9 cases by culture of the excised valve; in 3 by microscopy of the excised valve, on which large numbers of Gram positive cocci were seen although the culture was sterile; and in the other 4 by isolation from a site other than the excised valve (2 respiratory specimens, 1 from the pacemaker tip, and 1 from an excised embolus). In addition 5 of the 6 cases of *Bartonella* infection were confirmed by polymerase chain reaction study of the excised valve. Two thirds of the 32 patients for whom no pathogen was identified had received antibiotics before blood was cultured. Thus truly "negative" endocarditis was very uncommon (6% of the cases).

Conclusion: If blood cultures are negative in definite or suspected endocarditis, serum should be analysed for *Bartonella*, *Coxiella*, and *Chlamydia* species antibodies, and the excised valve or (rarely) embolus should be analysed by microscopy, culture, histology, and relevant polymerase chain reaction. Other specimens may be relevant. The Duke criteria performed poorly in BCNE; St Thomas' additional minor criteria gave more definite diagnoses.

See end of article for authors' affiliations

Correspondence to:
Professor Susannah J Eykyn, Department of Microbiology, 5th Floor, North Wing, St Thomas' Hospital, Lambeth Road, London SE1 7EH, UK; susannah.eykyn@doctors.net.uk

Accepted
20 November 2002

Blood culture negative endocarditis (BCNE) was recognised by Osler at the beginning of last century.¹ The three last published series of BCNE date from 1979² when organisms such as *Coxiella* and *Bartonella* species were not recognised causes of BCNE, from 1995,³ where diagnostic Duke criteria⁴ were not fully discussed nor *Bartonella* mentioned, and from 2001,⁵ where *Bartonella* investigation is not mentioned. We present 63 cases with reference to these new aetiological agents and assess the 1994 Duke criteria, their later modifications,⁶ and our 1997 proposed modifications⁷ in BCNE.

METHODS

Data on all episodes of endocarditis with negative blood cultures seen at St Thomas' Hospital between 1975 and 2000 were collected prospectively and analysed retrospectively. The demographic aspects of the series, clinical, microbiological, and echocardiographic data, as well as diagnostic criteria and outcome, were studied. Patients were assessed by the clinical criteria defined by Durack and colleagues⁴ (box 1) and by proposed modifications (St Thomas) (box 2). Additional modifications by Duke⁶ are summarised and are discussed in the analysis of the 34 pathologically proven cases of native valve endocarditis (NVE). Results are expressed as mean (SD) when appropriate or as percentages. Statistical analyses were performed by use of SPSS for Windows (SPSS Inc, Chicago, Illinois, USA).

In 2000, Li and colleagues⁶ proposed the following modifications to the Duke criteria:

- Elimination of echocardiographic minor criteria caused by the widespread use of transoesophageal echocardiography (TOE)
- Considering bacteraemia caused by *Staphylococcus aureus* a major criterion regardless of whether the infection is nosocomially acquired or a removable source of infection is present
- Recommending that Q fever serology be a major criterion.

RESULTS

There were 48 cases of blood culture negative NVE, involving 31 male patients and 17 female patients. Their mean age was 54 (17) years. Twenty nine patients were referred from other hospitals (60%). Eleven male and four female patients had prosthetic valve endocarditis (PVE) with a mean age of 53 (18) years. Eight (53%) were referred from other hospitals. Forty three of the 63 patients presented between 1990 and 2000.

Affected valves

The mitral valve was affected in 13 of 48 patients with NVE (27%), the aortic in 22 (46%), and both in 6 (13%). The tricuspid valve was affected in six patients (13%) and one had aortic, mitral, and tricuspid involvement (2%). In PVE, the mitral

Abbreviations: BCNE, blood culture negative endocarditis; NVE, native valve endocarditis; PCR, polymerase chain reaction; PVE, prosthetic valve endocarditis; TOE, transoesophageal echocardiography; TTE, transthoracic echocardiography

Box 1: Duke criteria for diagnosis of infective endocarditis⁴**Definite infective endocarditis**

- Pathological criteria
 - microorganisms shown by culture or histology in a vegetation, or in a vegetation that has embolised, or in an intracardiac abscess or
 - pathological lesions: vegetation or intracardiac abscess present, confirmed by histology showing active endocarditis
- Clinical criteria, using specific definitions*: two major criteria, or one major and three minor criteria, or five minor criteria

Possible infective endocarditis

- Findings consistent with infective endocarditis that fall short of "definite" but are not "rejected"

Rejected

- Firm alternative diagnosis for manifestations of endocarditis, or
- Resolution of manifestations of endocarditis with antibiotics for four days or less, or
- No pathological evidence of infective endocarditis at surgery or necropsy, after administration of antibiotics for four days or less

Specific definitions of proposed criteria*Major criteria:**

- Positive blood culture for infective endocarditis:
 - typical microorganisms for infective endocarditis from two separate blood cultures (viridans streptococci, *Streptococcus bovis*, *Haemophilus* sp, *Actinobacillus* sp, *Cardiobacterium* sp, *Eikenella* sp, *Kingella* sp, or community acquired *Staphylococcus aureus* or enterococci) in the absence or a primary focus, or
 - persistently positive blood culture, defined as recovery of a microorganism consistent with infective endocarditis from (a) blood cultures drawn more than 12 hours apart; or (b) all of three or a majority of four or more separate blood cultures, with first and last drawn at least one hour apart
- Evidence of endocardial involvement
 - positive echocardiogram for infective endocarditis: (a) oscillating intracardiac mass, on valve or supporting structures, or in the path or regurgitant jets, or on implanted material, in the absence of an alternative anatomic explanation; or (b) abscess; or (c) new partial dehiscence of prosthetic valve or
 - new valvar regurgitation

Minor criteria

- Predisposition: predisposing heart condition or intravenous drug use
- Fever $\geq 38^{\circ}\text{C}$
- Vascular phenomena: major arterial emboli, septic pulmonary infarcts, mycotic aneurysm, intracranial haemorrhage, conjunctival haemorrhages, or Janeway lesions
- Immunological phenomena: glomerulonephritis, Osler's nodes, Roth's spots, or rheumatoid factor
- Microbiological evidence: positive blood culture but not meeting a major criterion or serological evidence of active infection with organism consistent with infective endocarditis
- Echocardiogram consistent with infective endocarditis but not meeting a major criterion

valve was involved in six cases (40%), the aortic in eight (53%) and both in one (7%). Seven patients had early PVE (less than one year from surgery) and eight had late PVE.

Organisms

Blood cultures were negative in all 63 cases but in only 32 (51%) was no causative organism found. Table 1 shows organisms and diagnostic studies. All *Bartonella* species detected by serology were also grown or identified by polymerase chain

Box 2: Proposed additional minor criteria for the Duke classification⁷

- Newly diagnosed splenomegaly
- Newly diagnosed clubbing
- Splinter haemorrhages
- Petechiae
- High erythrocyte sedimentation rate, defined as more than 1.5 times the upper limit of normal (> 30 mm/h for patients < 60 years of age; > 50 mm/h for those > 60 years of age)
- High C reactive protein concentration, defined as > 100 mg/l
- Microscopic haematuria*
- Central non-feeding venous lines
- Peripheral venous lines

*Haematuria was disregarded for patients with positive urine cultures, menstruating women, patients with end stage renal disease, and patients with urinary catheters.

Table 1 Causative organism and diagnostic investigations

Diagnostic test	Native valve	Prosthetic valve	Total
Serology			
<i>Bartonella</i> sp	1		1
<i>Chlamydia psittaci</i>	1		1
<i>Coxiella burnetii</i>	5	3	8
Serology + PCR (<i>Bartonella</i> sp)	5		5
Valve culture	6*	3†	9
Culture of other material	4‡		4
Valve microscopy	2§	1¶	3
No organism	24	8	32

**Streptococcus bovis*, *Streptococcus mitis*, *Enterococcus faecium*, *Histoplasma capsulatum*, *Aspergillus* sp, *Aspergillus fumigatus*.

†*Rhodococcus* sp, *Staphylococcus aureus*, *Staphylococcus epidermidis*.

‡*S aureus* (sputum 1, pleural fluid 1), coagulase negative staphylococcus (pacemaker tip), viridans streptococcus (embolus).

§Gram positive coccus.

¶Yeast.

reaction (PCR) of valve material from 1995 onwards. Twenty two of the 32 (69%) patients for whom no organisms were identified had been given antibiotics before blood samples were drawn.

Predisposing conditions

Table 2 shows cardiac and non-cardiac conditions known to predispose to endocarditis. Six of 36 patients (17%) who had Gram positive cocci seen or grown in sites other than blood ($n = 4$) or did not have any identifiable organism ($n = 32$) had poor dentition or recent dental work. Two of the five patients with *Coxiella burnetii* endocarditis lived on farms and one frequently ate goat's milk and soft cheeses. The patient with *Chlamydia psittaci* endocarditis kept racing pigeons. The patient with *Bartonella henselae* endocarditis kept cats that had recently had fleas. The patient with *Bartonella berkhoffii* had contact with dogs, pigs, and a cockerel. Two of the three patients with *Bartonella quintana* "lived rough"; the third did not but kept birds.

Clinical features

Table 3 presents the main clinical features. The duration of symptoms was 7 (6) weeks for the 34 cases of known NVE and 7 (4) weeks in the 12 cases of PVE.

Echocardiography

Forty one of the 48 patients (85%) with NVE had transthoracic echocardiograms (TTE) and 8 (17%) had TOE. TTE showed major criteria for endocarditis in 32 patients (78%) and minor

Table 2 Conditions predisposing to blood culture negative endocarditis

Condition	Number (%)
Cardiac	
Prosthetic valve	13 (24)
Prosthetic valve and previous IE	2 (4)
Bicuspid aortic valve	7 (15)
RHD	4 (8)
Aortic stenosis	3 (6)
Calcified aortic and mitral valves	1 (2)
Ventricular septal defect	1 (2)
Mitral stenosis	1 (2)
RHD and previous IE	1 (2)
Mitral valve prolapse	1 (2)
Non-cardiac	
Pacemaker	3 (6)
Intravenous drug use	2 (4)
Atrioventricular fistula	1 (2)
None	23 (48)

IE, infective endocarditis; RHD, rheumatic heart disease.

criteria in 8 (20%) and was non-contributory in just one case. TOE showed major criteria in two cases where TTE was unhelpful. Thirteen of the 15 patients with PVE had TOE (87%); of these, 10 (77%) met major criteria for endocarditis (four had abscess). One TOE showed only a minor criterion and two were non-contributory.

Complications

Table 4 lists the complications.

Surgery and mortality

Thirty six of the 48 patients with NVE had surgery (75%), of whom 33 survived (92%). Three of the six patients who did not have surgery died. Causes of death in these patients were pancreatic carcinoma, sudden arrest on admission, and heart failure. Two of the three patients who had surgery and died had unrelated causes of death and one died in the operating theatre. Overall mortality in BCNE affecting native valves was 6 of 48 (12.5%) and mortality directly related to surgery was 1 of 36 (3%).

Thirteen of the 15 patients with BCNE affecting prosthetic valves had surgery (87%), of whom two died (15%) of cardiac causes. Both had early PVE (one and two months, respectively) and had emergency valve replacements (one aortic valve and one mitral valve). Two patients with PVE did not have surgery and survived: one was 43 years old, had had a

Table 3 Clinical features of blood culture negative endocarditis

Feature	Native valve (n=48)	Prosthetic valve (n=15)
Fever		
≥38–<39°C	24 (50%)	6 (40%)
≥39°C	9 (19%)	3 (20%)
Absent	8 (17%)	3 (20%)
Not known	7 (15%)	3 (20%)
Cerebral emboli	10 (21%)	2 (13%)
Splinter haemorrhages	9 (19%)	3 (20%)
Haematuria	8 (17%)	4 (27%)
Splenomegaly	7 (15%)	4 (27%)
"Rash"	6 (13%)	
Clubbing	5 (10%)	
Pulmonary emboli	4 (8%)	
Peripheral emboli	2 (4%)	1 (7%)
Subconjunctival haemorrhage	2 (4%)	
Osler's nodes	2 (4%)	
No immunological/vascular features	14 (29%)	

Table 4 Complications in blood culture negative endocarditis

Complication	Native valve (n=48)	Prosthetic valve (n=15)
Heart failure*	24 (50%)	10 (67%)
Cardiovascular accident†	7 (15%)	2 (13%)
Valvar abscess	2 (4%)	2 (13%)
Renal failure	2 (4%)	
Pulmonary‡	3 (6%)	
Coronary artery aneurysm	1 (2%)	
Peripheral emboli	1 (2%)	1 (2%)
Complete heart block	1 (2%)	
None	11 (23%)	1 (7%)

*Includes acute valvar regurgitation with left ventricular failure.

†One case of cerebellar haemorrhage.

‡Infarct, abscess, or empyema.

prosthetic aortic homograft for seven months, and was cured after two weeks of receiving benzylpenicillin plus gentamicin; the other was 81, had had an aortic prosthesis for nine years, and had *C burnetii* endocarditis. He received doxycycline and rifampin for many months and died of unrelated causes. His antibody titre was still positive when last checked. Surgical mortality in the PVE group was 2 of 13 (15%) and overall mortality 2 of 15 (13%).

Analysis of diagnostic criteria in 34 cases of pathologically proven BCNE affecting native valves

Diagnosis of infective endocarditis was pathologically proven in 34 cases: 19 by macroscopic surgical findings, 13 by excised valve histology, and 2 by histopathology of valves at post-mortem examination.

The clinical diagnosis of endocarditis was definite according to the Duke criteria in 7 (21%) of the 34 cases, and all were based on one major and three minor criteria. Our modifications of the Duke criteria⁷ upgraded probable cases to definite by adding one or more minor criteria to the 14 cases with one major and two minor criteria. In 7 only one criterion was added (erythrocyte sedimentation rate or C reactive protein), in 1 case two criteria were added, in 5 cases three or more criteria were added, and in 1 case four minor criteria were added. These minor criteria were splenomegaly (3), haematuria (4), clubbing (2), splinters (2), rash (1), petechiae (2), and purpura (1) (table 5).

If we consider Duke's modifications of their criteria,⁶ 11 of 34 cases would be definite (32%). These 4 new definite cases would be based on considering *Coxiella* serology a major criterion. All 4 of these cases were upgraded by St Thomas' modifications based on minor criteria.

DISCUSSION

These 63 cases of BCNE constituted 12.2% of all cases of endocarditis seen in St Thomas' Hospital between 1975 and 2000. This accords with reported rates of between 3–23%.^{8–16} However, in only 6% of our 63 cases did we fail to detect a

Table 5 Assessment of diagnostic criteria in 34 cases of pathologically proven blood culture negative endocarditis affecting native valves

	Duke criteria	St Thomas' criteria
Definite endocarditis	7 (21%)	21 (62%)
Two major criteria*	0	0
One major ≥3 minor criteria	7	21
Probable endocarditis	27	11

*Blood culture positivity is a major criterion.

causative organism, which suggests that truly negative infective endocarditis is an uncommon diagnosis. In common with other series²⁻¹⁵ two thirds of our patients in whom Gram positive cocci were seen or grown from sites other than blood (valves, sputum, embolus, or pacemaker wire tip) had received antibiotics, suggesting a staphylococcal or streptococcal cause. Serology to detect antibodies to Gram positive cell walls and broad range PCR on valve material were not performed, but these investigations might have confirmed the identity of the causative pathogen.¹⁷⁻²¹ PCR amplification of specific gene targets and universal rRNA loci for bacteria and fungi has been done in blood, as well as in valve material from patients with endocarditis,²⁰ and might have contributed to diagnoses in our series. No studies were done for *Tropheryma whippelii*, another well described cause of BCNE.²²

The administration of antibiotics, especially aminoglycosides, also influences isolation of *Bartonella* species from blood and valve cultures,²³ but our cases of *Bartonella* endocarditis were diagnosed by serology or PCR of the excised valve, which have higher sensitivity. Nearly one quarter of our cases of BCNE were caused by "atypical" bacteria: *C burnetii* (8 of 63), *Bartonella* species (6 of 63) and *C psittaci* (1 of 3). This is in contrast to the last published series of BCNE where no positive serological results were seen for *Coxiella*, *Legionella* or *Chlamydia* species (*Bartonella* serology was not done).² Although cross reaction of *Bartonella* antibodies with *Chlamydia* antibodies is described,²⁴ our patient with *Chlamydia* antibodies had a strong epidemiological link to *Chlamydia* organisms, as he kept racing pigeons. *C burnetii* and *Bartonella* species are well recognised causes of BCNE,²⁵⁻³¹ and it is estimated that *Bartonella* species cause 3% of all cases of infective endocarditis. Four of our 63 cases (6%) were caused by fungi ("yeast" 1, *Aspergillus* species 2, *Histoplasma capsulatum* 1). This last case has been published.³² The sensitivity of blood culture in fungal endocarditis was 54% in a review of 270 cases of fungal endocarditis, but fungi caused less than 10% of cases of endocarditis overall.³³

Prosthetic valves and bicuspid aortic valves were the most frequently infected in our series (24% and 15%, respectively), which is in contrast to the series reported by Pesanti,² with 52 cases of BCNE, in which 50% of the patients had rheumatic heart disease and only 8% had prosthetic valves. Bicuspid aortic valves were not specifically mentioned but they were the most common cardiac locations of NVE in two recent series.^{7,11,34}

Seventeen per cent of patients did not have fever, which is a higher incidence than usually reported with positive blood cultures^{6-8,11} but is similar to that of 19% reported by Hoen and colleagues.³ This may be the result of previous administration of antibiotics. Patients with *Coxiella* and *Bartonella* endocarditis are likely to present with low grade or no fever.²⁵⁻²⁸ The complications and mortality are similar to recently published series of blood culture positive NVE and PVE.⁷⁻¹² However, when comparing our series with a recently published series of BCNE,⁵ our rate of severe complications such as heart failure and valvar regurgitation was less (50% v 78%); the rates of major embolic events (cerebral and pulmonary) were similar (29% v 34.4%). The sensitivity of echocardiograms for the diagnosis of endocarditis is similar to that of positive blood cultures.^{6-8,11} TTE showed a major criterion for endocarditis in 78% of our cases involving native valves; few of these patients had TOE. This is in contrast with our cases of BCNE involving prosthetic valves, where 13 of 15 patients had TOE, which showed major criteria in 78%. TOE is superior to TTE in diagnosing endocarditis in the subgroup of patients with negative blood cultures, as has been shown by Kupferwasser and colleagues⁵ studying 32 cases of native valve BCNE, and we agree it ideally should be performed in all of these patients.

Analysis of patients with negative blood culture by the Duke criteria shows that these criteria perform poorly in such patients, which is not surprising since blood culture positivity is a major criterion. This was also shown by Habib and colleagues¹⁵ where 22 of 93 patients with pathologically proven

infective endocarditis were misclassified as possible by the Duke criteria. Twenty one of the 22 patients in this series had negative blood cultures and 19 had one major and two minor criteria. In our series, the Duke criteria diagnosed only 21% of the pathologically proven native valve cases as definite, whereas the St Thomas' modifications diagnosed 64% ($p < 0.05$). This was achieved by the addition of one or more minor diagnostic criteria. It has been previously proposed that serology for *Coxiella* species be considered a major criterion for infective endocarditis.¹⁴ If we use Duke's modifications of the Duke criteria⁶ we would identify only 32% of pathologically proven native valve blood culture negative cases in the present series. It is our experience and that of others (Raoult D, 9th International Congress on Infectious Diseases, Buenos Aires) that rapidly developing clubbing and splenomegaly are common findings in patients with *Coxiella* and *Bartonella* deep seated infections such as endocarditis, and these features should be considered in the diagnostic criteria for endocarditis (as they were in the Beth Israel criteria). It may be argued that the Duke criteria would have performed better in the present study if TOE had been performed in all the patients, but as has already been pointed out, TOE was done in most PVE cases (13 of 15). TTE performed well in the NVE subset (78% major criteria detected).

In summary, BCNE may be a diagnostic dilemma but in many cases the causative pathogen can be determined. It is important to elicit a history of any previous antibiotic that the patient has taken; serum should always be analysed for antibodies to organisms that cannot be cultured by routine methods such as *C burnetii*, *Bartonella*, and *Chlamydia* species, and if available for antibodies to Gram positive bacterial cell walls; and excised valves or vegetations or other relevant material should undergo microscopy, culture, histopathology, and relevant PCR analysis. Lastly, it should be noted that the Duke criteria and their modifications are unsatisfactory because they yield few definite clinical diagnoses. They require amendments to accept serology for organisms besides *Coxiella* species as a major criterion and they should incorporate additional minor criteria, as has already been suggested.

Authors' affiliations

C C Lamas, S J Eykyn, Department of Infection, North Wing, 5th floor, St Thomas' Hospital, Lambeth Palace Road, London SE1 7EH, UK

REFERENCES

- 1 Osler W. Chronic infectious endocarditis. *QJM* 1909;**2**:219.
- 2 Pesanti EL, Smith IM. Infective endocarditis with negative blood cultures: an analysis of 52 cases. *Am J Med* 1979;**66**:43-50.
- 3 Hoen B, Selson-Suty C, Lacassin F, et al. Infective endocarditis in patients with negative blood cultures: analysis of 88 cases from a one-year nationwide survey in France. *Clin Infect Dis* 1995;**20**:501-6.
- 4 Durack DT, Lukes AS, Bright DK, and the Duke Endocarditis Service. New diagnostic criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. *Am J Med* 1994;**96**:200-9.
- 5 Kupferwasser LI, Darius H, Muller AM, et al. Diagnosis of culture-negative endocarditis: the role of the Duke criteria and the impact of transesophageal echocardiography. *Am Heart J* 2001;**142**:146-52.
- 6 Li JS, Sexton DJ, Mick N, et al. Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis. *Clin Infect Dis* 2000;**30**:633-8.
- 7 Lamas CC, Eykyn SJ. Suggested modifications to the Duke criteria for the clinical diagnosis of native valve and prosthetic valve endocarditis: analysis of 118 pathologically proven cases. *Clin Infect Dis* 1997;**25**:713-9.
- 8 Bouvet A. Human endocarditis due to nutritionally variant streptococci: *Streptococcus adjacens* and *Streptococcus defectivus*. *Eur Heart J* 1995;**16**(suppl B):24-7.
- 9 Sandre RM, Shaffran SD. Infective endocarditis: review of 135 cases over 9 years. *Clin Infect Dis* 1996;**22**:276-86.
- 10 Hogevik H, Olaison L, Andersson R, et al. Epidemiologic aspects of infective endocarditis in an urban population: a 5 year prospective study. *Medicine (Baltimore)* 1995;**74**:324-39.
- 11 Choudhury R, Grover A, Varma J, et al. Active infective endocarditis observed in an Indian hospital 1981-1991. *Am J Cardiol* 1992;**70**:1453-8.
- 12 Hoen B, Vandenesch F, Le Moing V, et al. Changing profile of infective endocarditis: results of a one year survey in France [abstract]. 6th International Symposium on Modern Concepts in Endocarditis and Cardiovascular Infections, Barcelona, 2001.

- 13 **Van Scoy RE**. Culture-negative endocarditis. *Mayo Clin Proc* 1982;**57**:149–54.
- 14 **Tunkel AR**, Kaye D. Endocarditis with negative blood cultures. *N Engl J Med* 1992;**326**:1215–7.
- 15 **Habib A**, Derumeaux. Value and limitations of the Duke criteria for the diagnosis of infective endocarditis. *J Am Coll Cardiol* 1999;**33**:2023–9.
- 16 **Kurland S**, Enghoff E, Landelius J, *et al*. A 10 year retrospective study of infective endocarditis at a university hospital with special regard to the timing of surgical evaluation in *S viridans* endocarditis. *Scand J Infect Dis* 1999;**31**:87–91.
- 17 **Bosshard PP**, Kronenberg A, Zbinden R, *et al*. Etiologic diagnosis of infective endocarditis by broad-range PCR and sequencing [abstract]. 6th International Symposium on Modern Concepts in Endocarditis and Cardiovascular Infections, Barcelona, 2001.
- 18 **Lang S**, Connaughton M, Lambert PA, *et al*. Rapid serodiagnosis of Gram-positive bacterial endocarditis [abstract]. 6th International Symposium on Modern Concepts in Endocarditis and Cardiovascular Infections, Barcelona, 2001.
- 19 **Connaughton M**, Lang S, Tebby SE, *et al*. Rapid serodiagnosis of Gram-positive endocarditis. *J Infect* 2001;**42**:140–4.
- 20 **Burnie JP**, Clark I. Immunoblotting in the diagnosis of culture negative endocarditis caused by streptococci and enterococci. *J Clin Pathol* 1995;**48**:1130–6.
- 21 **Millar B**, Moore J, Mallon P, *et al*. Molecular diagnosis of infective endocarditis: a new Duke's criterion. *Scand J Infect Dis* 2001;**33**:673–80.
- 22 **Fenollar F**, Lepidi H, Raoult D. Whipple's endocarditis: review of the literature and comparisons with Q fever, Bartonella infection, and blood-culture positive endocarditis. *Clin Infect Dis* 2001;**33**:1309–16.
- 23 **La Scola B**, Raoult D. Culture of Bartonella quintana and Bartonella henselae from human samples: a 5 year experience (1993 to 1998). *J Clin Microbiol* 1999;**37**:1899–905.
- 24 **Drancourt M**, Mainardi J, Brouqui P, *et al*. Bartonella (Rochalimaea) quintana endocarditis in three homeless men. *N Engl J Med* 1995;**332**:419–23.
- 25 **Harrison TG**, Doshi N. Serological evidence of Bartonella spp infection in the UK. *Epidemiol Infect* 1999;**123**:233–40.
- 26 **Raoult D**. Afebrile blood culture-negative endocarditis. *Ann Intern Med* 1999;**131**:144–5.
- 27 **Raoult D**, Fournier PE, Drancourt M, *et al*. Diagnosis of 22 new cases of Bartonella endocarditis. *Ann Intern Med* 1996;**125**:646–52.
- 28 **Roux V**, Eykyn SJ, Wyllie S, *et al*. Bartonella vinsonii subsp berkhoffii as an agent of afebrile blood culture negative endocarditis in a human. *J Clin Microbiol* 2000;**38**:1698–700.
- 29 **Musso D**, Raoult D. Coxiella burnetii blood cultures from acute and chronic Q-fever patients. *J Clin Microbiol* 1995;**33**:3129–32.
- 30 **Breathnach AS**, Hoare JM, Eykyn SJ. Culture-negative endocarditis: contribution of Bartonella infections. *Heart* 1997;**77**:474–6.
- 31 **Maurin M**, Birtles R, Raoult D. Current knowledge of Bartonella species. *Eur J Clin Microbiol Infect Dis* 1997;**16**:487–506.
- 32 **Wilmshurst PT**, Venn GE, Eykyn SJ. Histoplasma endocarditis on a stenosed aortic valve presenting as dysphagia and weight loss. *Br Heart J* 1993;**70**:565–7.
- 33 **Ellis ME**, Al-Abdely H, Sandrigde A, *et al*. Fungal endocarditis: evidence in the world literature, 1965–1995. *Clin Infect Dis* 2001;**32**:50–62.
- 34 **Lamas CC**, Eykyn SJ. Bicuspid aortic valve: a silent danger: analysis of 50 cases of infective endocarditis. *Clin Infect Dis* 2000;**30**:336–41.

IMAGES IN CARDIOLOGY


Left atrial thrombus caused by the primary antiphospholipid syndrome causing critical functional mitral stenosis

Intracardiac masses are rare and their exact nature is often difficult to ascertain by conventional imaging methods alone. A 40 year old white woman presented with a six month history of progressive dyspnoea, orthopnoea, and peripheral oedema. Clinically she had signs of biventricular failure. Images from her transthoracic echocardiogram showed a large left atrial mass causing critical functional mitral stenosis (below left and right). A differential diagnosis of vegetation, thrombus, myxoma or other cardiac tumour (for example, lymphoma) was considered. A thrombophilia screen demonstrated raised serum concentrations of both anticardiolipin antibody (aCL) and antibody to its cofactor, β_2 glycoprotein I (β_2 GPI) on two separate occasions. An autoimmune profile (including antinuclear antibody and antibodies to extractable nuclear antigens) was negative,


confirming the diagnosis of primary antiphospholipid syndrome.

The patient was initially managed with high intensity anticoagulation, but failed to improve on clinical or echocardiographic criteria, and therefore surgical resection of the mass was performed. Histopathological analysis of the mass confirmed old and recent thrombus undergoing organisation from the base and infiltrated by reactive macrophages. This case highlights the need to screen all patients with intracardiac masses of unknown aetiology for prothrombotic disorders.

S Yusuf
B P Madden
C W Pumphrey
syusuf@sghms.ac.uk



Two dimensional echocardiography (parasternal long axis view) showing the presence of a large left atrial mass (arrowed) closely associated with and prolapsing through the mitral valve. AMVL, anterior mitral valve leaflet; LA, left atrium; LV, left ventricle; PMVL, posterior mitral valve leaflet.



Obstruction of the mitral valve orifice was dramatically demonstrated on M mode imaging and transmitral Doppler recordings were consistent with severe functional mitral stenosis (peak transmitral anterograde flow velocity 2.2 m/s). IVS, interventricular septum; LVPW, left ventricular posterior wall.