

# Proteomics - From Single Molecules to Biological Pathways

Sarah R. Langley<sup>1</sup>, Joseph Dwyer<sup>1</sup>, Ignat Drozdov<sup>1,2</sup>, Xiaoke Yin<sup>1</sup>, Manuel Mayr<sup>1</sup>

<sup>1</sup> King's British Heart Foundation Centre, King's College London, London, UK

<sup>2</sup> Centre for Bioinformatics - School of Physical Sciences and Engineering, King's College London, UK

**Correspondence to:** Prof. Manuel Mayr, King's British Heart Foundation Centre, King's College London, 125 Coldharbour Lane, London SE5 9NU, UK. Phone: +44 (0) 20 7848 5132, Fax: +44 (0) 20 7848 5296, email: manuel.mayr@kcl.ac.uk

Published on behalf of the European Society of Cardiology. All rights reserved.

© The Author 2012. For permissions please email: journals.permissions@oup.com

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that the original authorship is properly and fully attributed; the Journal, Learned Society and Oxford University Press are attributed as the original place of publication with correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oup.com

## Abstract

The conventional reductionist approach to cardiovascular research investigates individual candidate factors or linear signaling pathways but ignores more complex interactions in biological systems. The advent of molecular profiling technologies that focus on a global characterisation of whole complements, allows an exploration of the interconnectivity of pathways during pathophysiologically relevant processes, but has brought about the issue of statistical analysis and data integration. Proteins identified by differential expression as well as those in protein-protein interaction networks identified through experiments and through computational modelling techniques can be used as an initial starting point for functional analyses. In combination with other -omics technologies, such as transcriptomics and metabolomics, proteomics explores different aspects of disease and the different pillars of observations facilitate the data integration in disease-specific networks. Ultimately, a systems biology approach may advance our understanding of cardiovascular disease processes at a “biological pathway” instead of a “single molecule” level and accelerate progress towards disease-modifying interventions.

**Keyword:** Proteins – Metabolites – Mass spectrometry – Systems Biology - Bioinformatics

**This article is part of the Review Focus on: Cardiovascular Systems Biology**

## 1. Introduction

Proteomics represents the large-scale analysis of proteins, particularly their structures and functions. The term "proteomics" was coined to make an analogy with genomics, the study of the genome. While the genome is just the "blueprint" for the proteins, the proteins execute cellular function. Importantly, the transcriptome is not linearly proportional to proteome and many human diseases result from alterations in the proteome. In the first part of this review, we provide a short summary of proteomics techniques that have been extensively reviewed elsewhere<sup>1-3</sup>. Knowing the major limitations and advantages of the different proteomic techniques is essential for their successful application. An overview of systems biology approaches and examples follows, along with some of the resources available. Computational methods for dealing with the unique challenges of proteomics data will be key to fulfil the promise of systems biology.

## 2. Proteomics

Before summarizing different proteomic strategies (**Figure 1**), a few points should be emphasized<sup>4</sup>:

- 1) No proteomic technology can currently resolve the entire complexity of the mammalian proteome.
- 2) With any proteomic technique, there is bias towards more abundant proteins.
- 3) In general, there is a trade off between how many proteins can be quantified and how accurately they can be quantified.
- 4) Inevitably, information is lost by the propagation of quantitative peptide information to protein changes. **Table 1** gives a brief overview of the advantages and disadvantages of the following proteomics techniques.

**2.1. Two-dimensional gel electrophoresis (2-DE).** 2-DE gels allow separation of proteins based on their isoelectric point (pI) and molecular weight (Mw)<sup>5</sup>. The first dimension involves separating proteins according to their pI. A protein mixture is loaded onto a strip with an immobilised pH gradient. Once an electric field is applied, the proteins migrate to their pI where they become zwitterionic, i.e. they lose their net charge and stop migrating (isoelectric focusing). After isoelectric focusing is complete, the immobilised pH gradient strips are transferred onto large-format gels for separation in the second dimension, where proteins are resolved according to their molecular mass by SDS-PAGE.

Unlike SDS-PAGE, 2-DE gels produce complex maps of proteomes that are visualised as discrete protein 'spots'. Since pI and Mw are independent properties, 2-DE gels can resolve many more proteins than SDS-PAGE. Importantly, the same protein may be present in multiple spots on a gel. Shifts in pI or molecular weight indicate the presence of post-translational modifications, protein degradation or protein isoforms<sup>6, 7</sup>. Protein features are visualised with Coomassie or silver staining and differential expression between samples is determined using relative densitometric quantification. However, gel-to-gel variability can limit the quantitative accuracy and prohibit the detection of minor differences in expression.

A more sophisticated 2-DE technique is difference gel electrophoresis (DIGE, **Figure 2A**)<sup>8</sup>. DIGE involves fluorescent labelling of protein mixtures with Cy-dyes in order to determine relative differences in protein expression. An internal standard comprising of the pooled experimental samples is included, which is representative of all samples. The sensitivity of detection of DIGE is comparable with the sensitivity of silver staining<sup>9</sup> and the dyes are matched for pI and Mw. The main advantage of DIGE over

conventional 2-DE gels is that samples can be multiplexed on the same gel thus reducing the number of gels needed and limiting experimental variation. DIGE employed with an internal standard reliably quantifies differences as low as 10% in protein expression<sup>10</sup>. The gels are scanned using a fluorescence scanner, which specifically measures the emission wavelength of each Cy-dye. Commercial software packages match protein features and calculate differential expression from the scanned gel images. Normalisation of protein levels across gels is performed by comparing the protein ratios to the internal standard that is co-detected on each gel.

Unlike other proteomic techniques, quantitation by 2-DE is performed at the protein, not the peptide level and the quantitation is uncoupled from the identification by mass spectrometry (MS). Silver staining can be used to visualise protein features on a gel to facilitate excision of the relevant spots for MS. Alternatively, spots are directly picked from fluorescent gels using a robotic spot picker. Spots are then subjected to in-gel tryptic digestion before protein identification.

One of the main caveats of the 2-DE approach is that high abundant proteins mask less abundant proteins. This can be partially addressed by using gradients with a narrow pH range (**Figure 2B**). However, separation in the first dimension and in particular the transition from the first to the second dimension is not loss free and very large, small, and hydrophobic proteins remain difficult to resolve.

**2.2. Liquid-chromatography tandem mass spectrometry (LC-MS/MS).** LC-MS/MS is the current gold standard in proteomics. The basic principle of MS involves measuring the mass to charge ratio ( $m/z$ ) of an ionised peptide and its fragmentation products. Proteins are initially digested by enzymes such as trypsin to produce peptide fragments that are easier to resolve by reverse-phase LC and ionize by electrospray MS<sup>11</sup>.

Depending on their hydrophobicity, the peptides elute at different time points from the reverse phase column (retention time). A typical workflow using LC-MS/MS involves a regular survey scan to record the masses and the intensities of the eluting peptides. The most abundant precursor ions eluting from the column are selected for fragmentation (MS/MS). The amino acid sequence information obtained from MS/MS data allows the identification of the protein. Peptide parameters, such as spectral counts, ion intensities and chromatographic peak area, can provide a quantitative index for protein abundance (label-free quantitation)<sup>12</sup>. The versatility of mass spectrometric technology has spawned numerous different mass spectrometers with MALDI-TOF-TOF, Q-TOF and Orbitrap mass analysers<sup>13</sup> being among the common ones currently in use for discovery proteomics.

**2.3. Gel-LC-MS/MS.** Pre-fractionation by SDS-PAGE prior to MS has proven useful in the characterization of samples that are not amenable to separation by 2-DE. It also helps to overcome the single greatest cause of bias against low abundance proteins - the stochastic under sampling of low abundance peptides because high-abundance peptides dominate the duty cycle of the mass spectrometer. For gel-LC-MS/MS analysis, proteins are separated by SDS-PAGE, the entire gel lane is divided into a series of bands, the bands are excised without leaving empty gel pieces behind, digested with trypsin and LC-MS/MS analysis is performed on each of the bands<sup>14, 15</sup>. Since gel bands tend to be mixtures of proteins, LC separation is essential for protein identification and quantitation, i.e. by spectral counting<sup>16</sup>. Spectral counting has become a popular strategy to quantitate relative protein abundance but is less reliable for complex mixtures. Generally, the more abundant a protein, the more likely it is detected by MS/MS. The spectral counts are derived from the number of MS/MS spectra

corresponding to a particular protein.

In the gel-LC-MS/MS approach, information on the native Mw of a protein is preserved. If protein degradation has occurred prior to tryptic digestion, peptides are detected by MS in gel segments below the expected Mw of the native proteins (**Figure 2C**). Thus, verification whether differentially expressed proteins are confined to the same gel bands is essential. Otherwise, a degraded protein may appear upregulated due to its characteristic “laddering” on the SDS-PAGE (**Figure 2D**). Alternatively, protein fragments may be too small and escape detection because they migrated ahead of the buffer front. On the other hand, information on proteolytic degradation products is important and lost in conventional shotgun proteomics analysing tryptic peptides without prior separation at the protein level.

**2.4. Shotgun proteomics.** Apart from gel-based approaches, there are gel-free methods to quantify differences in protein expression based on peptide abundance. While these shot-gun proteomic methods can mine deeper into the proteome, problems arise with quantitation if samples are too complex. MS is not inherently quantitative because of differences in the ionization efficiency. The most abundant ions will attract the most charges during electrospray ionization, making it less likely for low-level peptides to get ionized. To avoid false positive protein changes due to co-eluting high-abundant peptides, labelling techniques should be used for reliable quantitation. Popular labelling methods include isobaric tagging for relative and absolute quantification (iTRAQ), tandem mass tags (TMT) and stable isotope labelling by amino acids in cell culture (SILAC) <sup>17</sup>. iTRAQ is currently available as 4-plex and 8-plex, allowing the relative quantification of up to eight samples, whereas labelling of TMT and SILAC can be used with six and three samples, respectively<sup>18</sup>. However, peptides are just a

surrogate measure and not always reliable for protein quantitation, i.e. if they are subject to post-translational modifications or proteolysis.

**2.4.1. SILAC.** SILAC makes use of non-radioactive isotope labels to label proteins with light (eg.  $^{12}\text{C}$ ) and heavy isotopes (eg.  $^{13}\text{C}$ )<sup>18</sup>. Samples can be multiplexed and analysed during the same MS run therefore minimising experimental error<sup>19</sup>. The SILAC pairs co-elute during chromatography but the corresponding peptides of the heavy and light isoform appear with a characteristic mass shift. The relative quantity of each protein can be calculated by the differences in the peak intensities of SILAC labelled peptides. The use of SILAC to quantify differential levels of proteins goes beyond using cells in culture. SILAC labelled mice have been described with near complete labelling of all proteins although the SILAC diet is expensive<sup>20</sup>. Metabolic labeling also introduces information on amino acid synthesis and sourcing, protein assembly and turnover kinetics.

**2.4.2. iTRAQ, TMT.** In instances where human tissue is used, iTRAQ or TMT are the only option for multiplexing clinical samples for differential expression studies by LC-MS/MS<sup>21</sup> but these techniques are not without caveats<sup>22</sup>: 1) One disadvantage of the iTRAQ and the TMT system over SILAC is the fact that labelling is performed at the peptide level and occurs late in the experimental process. Before labelling, proteins are first extracted from cells or tissues and digested to peptides. This is a potential source of variation. 2) Unlike SILAC, quantitation is performed at the MS/MS, not the MS level. The peptides from different samples maintain their identical mass to charge (m/z) ratios after labelling (MS). Only upon fragmentation (MS/MS), the isobaric mass tags release their different reporter ions with a single isotopic substitution per tag and provide quantitative information for each individual sample. A commonly observed problem in



iTRAQ experiments is that a complex background can lead to underestimation of protein fold changes. During precursor ion selection, more than one peptide may be within the mass window selected for fragmentation. In such mixed MS/MS spectra, reporter ions originating from peptides of different proteins are erroneously combined for quantification.

**2.5. Protein identification.** While accurate and accessible databases are needed for each of the 'omics' fields, proteomics is perhaps the most dependent on these resources. The technologies for identifying and quantifying proteins are reliant on comprehensive databases for protein identification and peptide quantification. Whilst not under the scope of systems biology, these databases provide a foundation for the latter analyses as the curation and maintenance of these databases is vital for the correct identification and quantification of the examined proteins.

For functional and sequence based databases, UniProt is one of the most comprehensive. UniProt consists of several classifications: Swiss-Prot and TrEMBL contain sequence and functional information about proteins, UniRef and UniParc contain sequence and archived sequence records and, when available, supporting data such as literature references and cross-referenced databases<sup>23</sup>. Programmes such as Mascot, SEQUEST or X!Tandem, search FASTA protein sequences obtained from public databases such as UniProt. After performing an "*in silico* fragmentation" with known enzyme specificity, the peak mass lists with intensities (the experimental data) are searched against the *in silico*-fragmented database. Parent ion masses are scanned against the masses derived from the database sequences. If there is a match within a certain mass tolerance, the observed MS/MS spectra are then compared with the theoretical sequence-derived ion series. While not explicitly covered here, the review

and commentary by Noble and MacCoss provide insight into these methodologies and techniques<sup>24</sup>. The scoring algorithms can produce different results and the reliance on single-peptide identifications in large-scale datasets is a potential cause of false identifications. Most proteomic studies only report identifications with a minimum of two unique peptides or include the MS/MS spectra for single-peptide identifications.

### 3. Systems biology approaches.

**3.1 Cellular and subcellular proteome identification.** Technological advances in the past 5-10 years have made large 'omics' experiments feasible, where biological changes can be assessed at the systems level rather than at an individual species level (**Figure 3**). High throughput proteomics is no different as one can now identify and quantify the proteins present in a specific cell type or subcellular fraction. As proteins can be present at varying levels in different cellular systems, it is imperative to know the baseline measurements for cells and systems specific to cardiovascular disease. In that respect, the proteomes of several cardiovascular specific cell types have been characterised in the past few years, including human arterial smooth muscle cells<sup>25</sup>, human early pro-angiogenic cells<sup>26</sup>, rat cardiac stem cells and neonatal cardiomyocytes<sup>27</sup> and human left ventricle<sup>28</sup>. Recently, Burkhardt et al. characterised the proteome of human platelets within and between healthy subjects. They identified approximately 4,000 unique proteins and showed that 85% of the platelet proteome did not vary across subjects<sup>29</sup>. Subcellular fractions can also be informative for cardiovascular disease. Several of these fractions have been analysed, including the extracellular matrix (ECM) in human aorta<sup>14</sup>, the mitochondrial proteome in mouse<sup>30</sup> and the rodent cardiac myofilament<sup>31</sup>.

As an example, **Figure 4** illustrates the proteomic network structure of ECM proteins in two different murine cell types: primary aortic smooth muscle cells (**Figure**

**4A)** and cardiac fibroblasts (**Figure 4B**). Both are correlation networks where the links between nodes (proteins) represent correlation values  $>0.9$ . The Venn diagrams show the total number of proteins identified in the two cell types and those in common between the two (**Figure 4C**); the number of nodes (proteins) in the networks as well as the number of shared nodes between the two (**Figure 4D**); finally, the number of links (correlations  $> 0.9$ ) for SMC and CF as well as the ones in common between the two (**Figure 4E**). The two networks and the corresponding nodes and links highlight the differences in the relationship between the ECM proteins in two different cell types.

For a systems biology approach of cardiovascular disease, it is important to identify and quantify proteomes in different species, tissue, cellular or subcellular compartments, as the differences, shown here in two ECM-producing cell types, may be specific to the defined system. The inclusion of these proteomes in public repositories will aid further systems biology as studies as the proteomics data will be available to other researchers. The biggest public repository of proteomics data is the PRoteomics IDentifications (PRIDE) database supported the European Bioinformatics Institute (EBI)<sup>32</sup>. As of the data of submission for this review, PRIDE contained over 26,000 proteomics experiments with the associated studies.

**3.2. Differential protein expression analysis.** With a defined system, one can study changes that result after a systematic perturbation. These perturbations can come in the form of inhibition, over-expression, incubation or a number of other cellular manipulation techniques, but also as a comparison between normal and disease samples. The system perturbation approach is not unique to proteomics; the other 'omics' fields use similar approaches; a large number of studies have been performed on transcriptomics and differential gene expression analysis and several methods have

been developed. Proteomics faces similar statistical and computational considerations genomics, but there are also challenges specific to proteomics, especially label-free techniques. Several studies, including our own<sup>14, 15, 33, 34</sup>, have applied standard statistical methods, to cardiovascular proteomics data<sup>35-37</sup>. While these methods may be appropriate, i.e. for analysis of DIGE data, they are not optimal for label free, spectral count protein expression data. Although many proteins are identified, less abundant proteins often contain one or more missing values across samples<sup>38</sup>. This presence/absence dichotomy is not suitable for basic imputation methods. Missing values will skew statistical tests that assume normality and standard statistical methods (Student's t-test, Analysis of Variance - ANOVA, linear regression) may not accurately determine differential expression<sup>39-41</sup>. On the other hand, excluding proteins with missing spectral counts will inherently create a bias towards high abundant proteins<sup>39</sup>. Low abundant proteins, however, are often informative, especially when comparing disease states, as the presence in one state and the absence in another can suggest a functional role for that protein.

The small number of replicates within an experiment reduces the robustness and increases the noise to signal ratio<sup>42</sup>. Non-parametric tests, like the Wilcoxon rank-sum test, have limited power when used with the small sample sizes often found in proteomic studies<sup>41</sup>. Applying multiple testing corrections becomes problematic as permutation or bootstrapping techniques are not feasible with small sample sizes. With this in mind, several methods have been developed specifically for evaluating protein differential expression, which take into consideration these limitations. Some methods address the non-normal distribution properties of the data, where the data is normalised and transformed to better fit the standard statistical tests. Three of the commonly used methodologies for this approach are the Normalized Spectral Abundance Factor (NSAF)

<sup>43</sup>, the Power Law Global Error Model (PLGEM) <sup>44</sup> and the Normalized Spectral Index (SI<sub>N</sub>)<sup>45</sup> (**Table 2**). These methods, however, do not take into consideration the small sample sizes that are common in proteomics experiments nor do they directly correct for multiple testing.

Other methods incorporate techniques that address both the non-normal distributions and the limited number of replications. The Spectral Index (Spl)<sup>46</sup>, Qspec<sup>47</sup>, and the hybrid approach proposed by Wang *et al.*<sup>39</sup>, are three examples of methods that account for small sample sizes and do not require the data to be normally distributed (**Table 2**). These methods also directly incorporate multiple testing corrections. Unlike gene expression microarray analyses, there is no standard method for normalisation and differential expression analyses in proteomics. Due to the variability between experiments and methodologies, statisticians and computational biologists should guide proteomic analyses.

**3.3. Incorporating functional and pathway information.** Functional information such as gene ontology (GO) and pathway resources can inform on the biological function of proteins and their interactions and on the relevance of the proteins to the disease. GO<sup>48</sup> and Kyoto Encyclopedia of Genes and Genomes (KEGG)<sup>49</sup> are the two resources widely used in the literature but there are other pathway and functional databases available (see **Table 3**). The KEGG database contains manually curated pathways within five areas: metabolism, genetic information processing, environmental information processing, cellular processes and human disease. Unlike GO, these pathways are species dependent. The GO database contains species independent terms relating to genes and their products. There are three main classifications of ontologies, cellular component (CC), biological process (BP) and molecular function with several sub-

classifications under each of the three. In addition, there is a special GO consortium specifically focused on annotating genes relevant to cardiovascular disease (<http://www.geneontology.org/GO.cardio.shtml>), and, to date, have identified over 4,000 genes with a cardiovascular disease association. The Go Cardiovascular Consortium also annotates gene products, including proteins and microRNAs. Instead of a one-way exchange, the relationship between the cardiovascular proteomics community, including our group, and the consortium is circular. Researchers not only use GO annotation to inform their research, but can also submit data from their experiments to validate annotations and suggest novel cardiovascular GO terms.

As an example, Isserlin *et al.*<sup>36</sup> incorporated a differential expression analysis with a Gene Set Enrichment Analysis (GSEA) to identify sets of differentially expressed proteins that were enriched for functional terms relating to dilated cardiomyopathy. They utilized GO as well as several other sources of publicly available functional data to perform the GSEA and derived functional networks, which show novel processes in the progression from pre-symptomatic to full dilated cardiomyopathy.

**3.4. Network biology.** The identification of differentially expressed proteins is only one part of a systems biology approach to proteomics. Further analyses are often performed on the set of differentially expressed proteins to elucidate their functional role in the disease pathology. In the conventional reductionist approach, experiments tend to address intracellular signalling cascades as linear models with the involved molecules confined to a signal pathway. The post-genomic shift in paradigm acknowledges the fact that many biological systems can be represented using concepts of network biology. Different pathways cross-talk with each other at points that can be graphically represented as well connected nodes or nexuses within a map of signalling networks<sup>50</sup>.

In addition to high-throughput data acquisition, the last decade introduced a number of sophisticated methodologies that intend to interrogate cellular interactions<sup>51-55</sup>. Preliminary analyses of these interactomes revealed the complexity of molecular signalling, which presents a challenge for accurate interpretation and application. It is now believed that the human interactome is comprised of ~20,000 protein-coding genes, ~1,000 metabolites, and an undefined number of distinct proteins, while the number of functional links between these components is expected to be ~130,000<sup>51</sup>. An emerging computational discipline, network biology, has been proposed as a tool that may supplement traditional quantitative analysis and uncover relational properties that control the behaviour of a cell through data integration and computational modeling<sup>56</sup>. Network biology was successfully used to define gene regulatory patterns in physiological cardiac hypertrophy<sup>57</sup> and highlight network topology of heart development and failure<sup>58</sup>. Increasing evidence suggest that combination of network concepts, such as centrality, with gene, protein or microRNA expression information may contribute to better prioritization of relevant biological targets<sup>59-61</sup>.

Despite their usefulness, networks analyses should be used with some degree of caution. It is currently not feasible to access and characterise the entire human proteome and so each proteomic network will consist of a subset of all possible proteins. Network studies, especially those focusing on protein-protein interactions, have shown that network properties from a sample or subset of a global network differ from the properties of the global network<sup>62-64</sup>. For example, biological networks have been described as 'scale-free' networks where there are several nodes with a high degree of connectivity and many nodes with a low degree of connectivity<sup>65</sup>. While this may be an appropriate assumption for a large-scale biological network, no statistical tests were applied to prove scale-freeness in biological networks and the smaller sub-networks,

including most protein-protein interaction networks, do not follow the same assumptions

#### **4. Is a systems-level integration of 'omics' data the way forward?**

Within a biological system, proteins do not act on their own, but rather, through complex interactions with metabolites, RNA, and other proteins. As we learn more about the pathophysiology of cardiovascular diseases, the underlying complexity becomes apparent, and the integration of 'omics' fields provides an unbiased way to elucidate the underlying mechanisms. The advancements in the technologies and the data availability of each of the 'omics' gives rise to a finer assessment but also provides a greater opportunity to study the complex interactions between genes, gene expression, proteins, and metabolites. Each of these 'omics' techniques provides a different level of information and by integrating them we can create new hypothesis and novel insights into disease.

**4.1. Integrating transcriptomics and proteomics.** Initial investigations into the correlation between mRNA levels and protein levels have shown poor to moderate associations between the two. These low correlations can be attributed to epigenetic factors, translation rates and protein degradation rates, but they can also be due to the levels being assessed different samples, across different time points<sup>66</sup>. To overcome some of these issues, Schwanhausser *et al* used mouse fibroblasts to quantify and analyse global mRNA and protein levels along with their associated half-lives, transcription and translation rates<sup>67</sup>. To get an accurate picture of the strength of association between mRNA and protein levels, the authors used metabolic pulse labelling in an experimentally growing population of embryonic fibroblasts to record mRNA and proteins levels occurring at the same time point. They found that 40% of the



variation in proteins levels can be attributed to variation in mRNA levels but translation efficiency was the best predictor. Certain combinations of half-lives and mRNA levels correspond to shared functional role, indicating shared selective pressures. In another transcriptomic and proteomic analysis, Zhao *et al.* reconstructed a heart specific metabolic network using transcriptome and proteome data with a model-building algorithm. Using generic genome-wide metabolic networks, they constructed heart specific models by mapping transcriptomic and proteomic data from the heart onto the genetic networks. The resulting model contained 2,803 reactions with 1,721 active enzymes in the heart. With this metabolic network, they were able to estimate the lethality, *in silico*, of house-keeping and heart specific genes and identify potential CVD biomarkers<sup>68</sup>.

**4.2. Integrating proteomics and metabolomics.** Currently, proteomics and metabolomics are rarely used in tandem, but this technological platform offers advantages: First, it has the potential to identify emergent behaviour that cannot be found by studying proteins or metabolites in isolation. Besides, proteomic and metabolomics findings can effectively reinforce or cross-validate each other. We utilized a combined proteomics and metabolomics approach to investigate cardiovascular diseases<sup>69</sup>. Our aim was to contribute to a better understanding of enzymatic and metabolite changes associated with cardiovascular diseases, including atherosclerosis<sup>6</sup>,<sup>70</sup>, ischemic preconditioning<sup>71</sup>, cardioprotective signalling<sup>72, 73</sup> and atrial fibrillation<sup>74</sup>. In a recent study on myocardial hibernation, murine hearts were analysed by a combined transcriptomic, proteomic and metabolomic approach (**Figure 5**)<sup>75, 76</sup>. Unguided network analysis correctly identified hypoxia-inducible factor 1 alpha (HIF1 $\alpha$ ) activation as the top signalling pathway, and provided independent confirmation that anaerobic glycolysis

is affected. A direct link to cardiac remodelling was also provided by the activation of collagen-hydroxylases, which produce hydroxyproline. By combining the 'omics' data, the p-value of the HIF1 $\alpha$  signalling pathway decreased by two orders of magnitude, and became the top-ranking pathway even though it was not the top-ranked pathway based on either dataset individually. The proteomics and transcriptomics focused on and contributed different molecules to the protein network, which enabled the HIF1 $\alpha$  signalling pathway to rise to the top-ranked pathway.

**4.3. Personal “-omics” profiles.** In a proof of concept study, Chen *et al.* presented an integrative personal -omics" profile (iPOP) analysis, where the genetic, transcripomic, proteomic, metabolomic and autoantibody profiles were measured and integrated for one healthy individual over the course of 14 months<sup>77</sup>. The measurements were assessed in blood components (plasma, serum and peripheral blood mononuclear cells (PBMC)) at several time points during the course of the study. The 'omic' responses were studied in greater detail during two viral infections, showing the dynamic response of the immune system. Interestingly, the authors identified a genetic predisposition to type II diabetes at the start of the study and noticed a pronounced change in insulin-related responses after the second infection. While the causal relationship between the infection and the onset of diabetes cannot be determined from one individual, these tightly linked events, and the indication that they are related, were only detected through the combination of 'omics' profiles. As the technology to measure 'omics' profiles becomes feasible, the greatest challenge will not be the generation of data, but their analysis.

## 5. Conclusions.

A discrete biological function is very rarely attributed to one single molecule; more often it is the combined input of many proteins. The studies mentioned above, which integrate protein data with other 'omics' data including transcriptomic<sup>67,68,77</sup>, metabolomic<sup>75,76,77</sup> and GWAS<sup>77</sup>, illustrate the utility of an integrative 'omics' approach to cardiovascular diseases. However, variants and changes from the genetic to the phenotypic level are not linearly associated and often variations seen at one level are absent at another. While the integration of data from different 'omics' techniques is still a challenge, the incorporation of proteomics with systems biology, and the application to study metabolism, is a promising area for future applications in cardiovascular diseases<sup>78, 79</sup>. Combining proteomics with stringent statistics, bioinformatics and other -omics technologies, such as metabolomics, can aid in identifying targets that have clinical relevance for working towards new therapies for cardiovascular disease<sup>80</sup>. Improvements in protein identification and quantification technologies as well as the availability of more proteomics datasets in public data repositories such as PRIDE<sup>32</sup>, combined with focused GO curation<sup>81, 82</sup> will facilitate the application of systems biology to cardiovascular research.

## Acknowledgement.

We are grateful to Drs. Basetti Madhu, Melanie Abbonenc and Athanasios Didangelos for providing Figures.

## Funding.

This work was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre at Guy's and St Thomas' NHS Foundation Trust and King's

College London. M.M. is a Senior Fellow of the British Heart Foundation.

**Conflict of Interest.** None.

## References

1. Prokopi M, Mayr M. Proteomics: A reality-check for putative stem cells. *Circ Res* 2011;**108**:499-511.
2. McGregor E, Dunn MJ. Proteomics of heart disease. *Hum Mol Genet* 2003;**12**:R135-144.
3. Arrell DK, Neverova I, Van Eyk JE. Cardiovascular proteomics: Evolution and potential. *Circ Res* 2001;**88**:763-773.
4. Mayr M, Zhang J, Greene AS, Gutterman D, Perloff J, Ping P. Proteomics-based development of biomarkers in cardiovascular disease: Mechanistic, clinical, and therapeutic insights. *Mol Cell Proteomics* 2006;**5**:1853-1864.
5. Gorg A, Weiss W, Dunn MJ. Current two-dimensional electrophoresis technology for proteomics. *Proteomics* 2004;**4**:3665-3685.
6. Mayr M, Chung YL, Mayr U, Yin X, Ly L, Troy H *et al.* Proteomic and metabolomic analyses of atherosclerotic vessels from apolipoprotein e-deficient mice reveal alterations in inflammation, oxidative stress, and energy metabolism. *Arterioscler Thromb Vasc Biol.* 2005;**25**:2135-2142.
7. McManus CA, Donoghue PM, Dunn MJ. A fluorescent codetection system for immunoblotting and proteomics through ecl-plex and cydye labeling. *Methods Mol Biol* 2009;**536**:515-526.
8. Unlu M, Morgan ME, Minden JS. Difference gel electrophoresis: A single gel method for detecting changes in protein extracts. *Electrophoresis* 1997;**18**:2071-2077.
9. Westermeier R, Scheibe B. Difference gel electrophoresis based on lys/cys tagging. *Methods Mol Biol* 2008;**424**:73-85.
10. McGregor E, Dunn MJ. Proteomics of the heart: Unraveling disease. *Circ Res*

- 2006;**98**:309-321.
11. Wilm M. Principles of electrospray ionization. *Mol Cell Proteomics* 2011;**10**:M111.009407.
  12. Liu H, Sadygov RG, Yates JR, 3rd. A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal Chem* 2004;**76**:4193-4201.
  13. Scigelova M, Hornshaw M, Giannakopoulos A, Makarov A. Fourier transform mass spectrometry. *Mol Cell Proteomics* 2011;**10**: M111.009431.
  14. Didangelos A, Yin X, Mandal K, Baumert M, Jahangiri M, Mayr M. Proteomics characterization of extracellular space components in the human aorta. *Mol Cell Proteomics* 2010;**9**:2048-2062.
  15. Didangelos A, Yin X, Mandal K, Saje A, Smith A, Xu Q *et al.* Extracellular matrix composition and remodeling in human abdominal aortic aneurysms: A proteomics approach. *Mol Cell Proteomics* 2011; **10**:M111.008128.
  16. Arnott D, Kishiyama A, Luis EA, Ludlum SG, Marsters JC, Jr., Stults JT. Selective detection of membrane proteins without antibodies: A mass spectrometric version of the western blot. *Mol Cell Proteomics* 2002;**1**:148-156.
  17. Bouyssie D, Gonzalez de Peredo A, Mouton E, Albigot R, Roussel L, Ortega N, *et al.* Mascot file parsing and quantification (mfpaq), a new software to parse, validate, and quantify proteomics data generated by icat and silac mass spectrometric analyses: Application to the proteomics study of membrane proteins from primary human endothelial cells. *Mol Cell Proteomics* 2007;**6**:1621-1637.
  18. Mann M. Functional and quantitative proteomics using silac. *Nat Rev Mol Cell Biol* 2006;**7**:952-958.

19. Ong SE, Mann M. A practical recipe for stable isotope labeling by amino acids in cell culture (SILAC). *Nat Protoc* 2006;**1**:2650-2660.
20. Kruger M, Moser M, Ussar S, Thievensen I, Lubber CA, Forner F *et al.* Silac mouse for quantitative proteomics uncovers kindlin-3 as an essential factor for red blood cell function. *Cell* 2008;**134**:353-364.
21. Dayon L, Sanchez JC. Relative protein quantification by ms/ms using the tandem mass tag technology. *Methods Mol Biol* 2012;**893**:115-127.
22. Karp NA, Huber W, Sadowski PG, Charles PD, Hester SV, Lilley KS. Addressing accuracy and precision issues in iTRAQ quantitation. *Mol Cell Proteomics* 2010;**9**:1885-1897.
23. Consortium U. Reorganizing the protein space at the universal protein resource (uniprot). *Nucleic Acids Res* 2012;**40**:D71-75.
24. Noble WS, MacCoss MJ. Computational and statistical analysis of protein mass spectrometry data. *PLoS computational biology* 2012;**8**:e1002296.
25. Dupont A, Corseaux D, Dekeyzer O, Drobecq H, Guihot A-L, Susen S, *et al.* The proteome and secretome of human arterial smooth muscle cells. *Proteomics* 2005;**5**:585–596.
26. Urbich C, De Souza AI, Rossig L, Yin X, Xing Q, Prokopi M, *et al.* Proteomic characterization of human early pro-angiogenic cells. *JMCC* 2011;**50**:333-336
27. Stastna M, Chimenti I, Marban E, Van Eyk JE. Identification and functionality of proteomes secreted by rat cardiac stem cells and neonatal cardiomyocytes. *Proteomics* 2010;**10**:245–253
28. Aye TT, Scholten A, Taouatas N, Varro A, Van Veen TAB, Vos MA *et al.* Proteome-wide protein concentrations in the human heart. *Mol. BioSyst* 2010;**6**:1917–1927

29. Burkhart JM, Vaudel M, Gambaryan S, Radau S, Walter U, Martens L, *et al.* The first comprehensive and quantitative analysis of human platelet protein composition allows the comparative analysis of structural and functional pathways. *Blood* 2012;**120**:e73-e82
30. Zhang J, Li X, Mueller M, Wang Y, Zong C, Deng N, *et al.* Systematic characterization of the murine mitochondrial proteome using functionally validated cardiac mitochondria. *Proteomics* 2008;**8**:1564–1575
31. Yin X, Cuello F, Mayr U, Hao Z, Hornshaw M, Ehler E, *et al.* Proteomics Analysis of the Cardiac Myofilament Subproteome Reveals Dynamic Alterations in Phosphatase Subunit Distribution. *Mol Cell Proteomics* 2010;**9**:497-509
32. Vizcaíno JA, Côté R, Reisinger F, Barsnes H, Foster JM, Rameseder *et al.* The Proteomics Identifications database: 2010 update. *Nucleic Acids Res* 2010;**38(Database issue)**:D736-42.
33. Mayr M, Siow R, Chung YL, Mayr U, Griffiths JR, Xu Q. Proteomic and metabolomic analysis of vascular smooth muscle cells: Role of pkcdelta. *Circ Res* 2004;**94**:e87-96.
34. Barallobre-Barreiro J, Didangelos A, Schoendube FA, Drozdov I, Yin X, Fernandez-Caggiano M *et al.* Proteomics analysis of cardiac extracellular matrix remodeling in a porcine model of ischemia/reperfusion injury. *Circulation* 2012;**125**:789-802.
35. Dai D-F, Hsieh EJ, Liu Y, Chen T, Beyer RP, Chin MT *et al.* Mitochondrial proteome remodelling in pressure overload-induced heart failure: The role of mitochondrial oxidative stress. *Cardiovasc Res* 2012;**93**:79-88.
36. Isserlin R, Merico D, Alikhani-Koupaei R, Gramolini A, Bader GD, Emili A. Pathway analysis of dilated cardiomyopathy using global proteomic profiling and



- enrichment maps. *Proteomics* 2010;**10**:1316-1327.
37. Hammer E, Goritzka M, Ameling S, Darm K, Steil L, Klingel K *et al.* Characterization of the human myocardial proteome in inflammatory dilated cardiomyopathy by label-free quantitative shotgun proteomics of heart biopsies. *J Proteome Res* 2011;**10**:2161-2171.
  38. Wang W, Zhou H, Lin H, Roy S, Shaler TA, Hill LR *et al.* Quantification of proteins and metabolites by mass spectrometry without isotopic labeling or spiked standards. *Anal Chem* 2003;**75**:4818-4826.
  39. Wang X, Anderson G, Smith RD, Dabney AR. A hybrid approach to protein differential expression in mass spectrometry-based proteomics. *Bioinformatics* 2012:1-8.
  40. Karp NA, Lilley KS. Design and analysis issues in quantitative proteomics studies. *Proteomics* 2007;**7** Suppl 1:42-50.
  41. Cairns DA. Statistical issues in quality control of proteomic analyses: Good experimental design and planning. *Proteomics* 2011;**11**:1037-1048.
  42. Käll L, Vitek O. Computational mass spectrometry-based proteomics. *PLoS Comp Biol* 2011;**7**:e1002277.
  43. Zybaylov B, Mosley AL, Sardu ME, Coleman MK, Florens L, Washburn MP. Statistical analysis of membrane proteome expression changes in *saccharomyces cerevisiae*. *J Proteome Res* 2006;**5**:2339-2347.
  44. Pavelka N, Fournier ML, Swanson SK, Pelizzola M, Ricciardi-Castagnoli P, Florens L *et al.* Statistical similarities between transcriptomics and quantitative shotgun proteomics data. *Mol Cell Proteomics* 2008;**7**:631-644.
  45. Griffin NM, Yu J, Long F, Oh P, Shore S, Li Y *et al.* Label-free, normalized quantification of complex mass spectrometry data for proteomic analysis. *Nat*

- Biotech* 2010;**28**:83-89.
46. Fu X, Gharib SA, Green PS, Aitken ML, Frazer DA, Park DR, Vaisar T *et al.* Spectral index for assessment of differential protein expression in shotgun proteomics. *J Proteome Res* 2008;**7**:845-854.
  47. Choi H, Fermin D, Nesvizhskii AI. Significance analysis of spectral count data in label-free shotgun proteomics. *Mol Cell Proteomics* 2008;**7**:2373-2385.
  48. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM *et al.* Gene ontology: Tool for the unification of biology. The gene ontology consortium. *Nat Genetics* 2000;**25**:25-29.
  49. Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. Kegg for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res* 2011;**40**:D109-D114.
  50. King JY, Ferrara R, Tabibiazar R, Spin JM, Chen MM, Kuchinsky A *et al.* Pathway analysis of coronary atherosclerosis. *Physiol Genomics* 2005;**23**:103-118
  51. Venkatesan K, Rual JF, Vazquez A, Stelzl U, Lemmens I, Hirozane-Kishikawa T *et al.* An empirical framework for binary interactome mapping. *Nat Methods* 2009;**6**:83-90.
  52. Rual JF, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, Li N *et al.* Towards a proteome-scale map of the human protein-protein interaction network. *Nature* 2005;**437**:1173-1178.
  53. Stelzl U, Worm U, Lalowski M, Haenig C, Brembeck FH, Goehler H *et al.* A human protein-protein interaction network: A resource for annotating the proteome. *Cell* 2005;**122**:957-968.
  54. Wu G, Feng X, Stein L. A human functional protein interaction network and its

- application to cancer data analysis. *Genome Biol* 2010;**11**:R53.
55. Ravasi T, Suzuki H, Cannistraci CV, Katayama S, Bajic VB, Tan K *et al*. An atlas of combinatorial transcriptional regulation in mouse and man. *Cell* 2010;**140**:744-752.
56. Vidal M, Cusick ME, Barabasi AL. Interactome networks and human disease. *Cell* 2011;**144**:986-998.
57. Drozdov I, Tsoka S, Ouzounis CA, Shah AM. Genome-wide expression patterns in physiological cardiac hypertrophy. *BMC Genomics* 2010;**11**:557.
58. Dewey FE, Perez MV, Wheeler MT, Watt C, Spin J, Langfelder P *et al*. Gene coexpression network topology of cardiac development, hypertrophy, and failure. *Circ Cardiovasc Genet* 2011;**4**:26-35.
59. Barrenas F, Chavali S, Holme P, Mobini R, Benson M. Network properties of complex human disease genes identified through genome-wide association studies. *PLoS One* 2009;**4**:e8090.
60. Zampetaki A, Kiechl S, Drozdov I, Willeit P, Mayr U, Prokopi M *et al*. Plasma microRNA profiling reveals loss of endothelial mir-126 and other microRNAs in type 2 diabetes. *Circ Res* 2010;**107**:810-817.
61. Zampetaki A, Willeit P, Tilling L, Drozdov I, Prokopi M, Renard JM *et al*. Prospective study on circulating microRNAs and risk of myocardial infarction. *J Am Coll Cardiol* 2012;**60**:290-299.
62. Stumpf MP, Wiuf C, May RM. Subnets of scale-free networks are not scale-free: Sampling properties of networks. *Proc Natl Acad Sci U S A* 2005;**102**:4221-4224.
63. Fernandes LP, Annibale A, Kleinjung J, Coolen ACC, Fraternali F. Protein networks reveal detection bias and species consistency when analysed by information-theoretic methods. *PloS one* 2010;**5**:e12083.

64. Annibale A, Coolen ACC. What you see is not what you get: How sampling affects macroscopic features of biological networks. *Interface Focus* 2011;**1**:836-856.
65. Stumpf MPH. From the cover: Subnets of scale-free networks are not scale-free: Sampling properties of networks. *Proc Natl Acad Sci USA* 2005;**102**:4221-4224.
66. de Sousa Abreu R, Pentalva LO, Marcotte EM, Vogel C, Global signatures of protein and mRNA expression levels. *Mol Biosyst* 2009;**5**,1512-1526
67. Schwanhausser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, *et al.* Global quantification of mammalian gene expression control. *Nature* 2011;**473**:337-342
68. Zhao Y, Huang J. Reconstruction and analysis of human heart-specific metabolic network based on transcriptome and proteome data. *BBRC* 2011;**415**:450-454
69. Mayr M. Metabolomics: Ready for the prime time? *Circ Cardiovasc Genet* 2008;**1**:58-65.
70. Mayr M, Grainger D, Mayr U, Leroyer AS, Leseche G, Sidibe A *et al.* Proteomics, metabolomics, and immunomics on microparticles derived from human atherosclerotic plaques. *Circ Cardiovasc Genet* 2009;**2**:379-388.
71. Mayr M, Metzler B, Chung YL, McGregor E, Mayr U, Troy H *et al.* Ischemic preconditioning exaggerates cardiac damage in PKC-delta null mice. *Am J Physiol Heart Circ Physiol* 2004;**287**:H946-H956.
72. Mayr M, Chung YL, Mayr U, McGregor E, Troy H, Baier G *et al.* Loss of PKC-delta alters cardiac metabolism. *Am J Physiol Heart Circ Physiol* 2004;**287**:H937-H945.
73. Mayr M, Liem D, Zhang J, Li X, Avliyakov NK, Yang JI *et al.* Proteomic and metabolomic analysis of cardioprotection: Interplay between protein kinase c epsilon and delta in regulating glucose metabolism of murine hearts. *J Mol Cell*

- Cardiol* 2009;**46**:268-277.
74. Mayr M, Yusuf S, Weir G, Chung YL, Mayr U, Yin X *et al.* Combined metabolomic and proteomic analysis of human atrial fibrillation. *J Am Coll Cardiol* 2008;**51**:585-594.
75. Mayr M, May D, Gordon O, Madhu B, Gilon D, Yin X *et al.* Metabolic homeostasis is maintained in myocardial hibernation by adaptive changes in the transcriptome and proteome. *J Mol Cell Cardiol* 2011;**50**:982-990.
76. May D, Gilon D, Djonov V, Itin A, Lazarus A, Gordon O *et al.* Transgenic system for conditional induction and rescue of chronic myocardial hibernation provides insights into genomic programs of hibernation. *Proc Natl Acad Sci U S A* 2008;**105**:282-287.
77. Chen R, Mias GI, Li-Pook-Tham J, Jiang L, Lam HYK, Chen R, *et al.* Personal Omics Profiling Reveals Dynamic Molecular and Medical Phenotypes. *Cell* 2012;**148**:1293–1307
78. Mayr M, Mayr U, Chung YL, Yin X, Griffiths JR, Xu Q. Vascular proteomics: Linking proteomic and metabolomic changes. *Proteomics* 2004;**4**:3751-3761.
79. Mayr M, Madhu B, Xu Q. Proteomics and metabolomics combined in cardiovascular research. *Trends Cardiovasc Med* 2007;**17**:43-48
80. Arrell DK, Terzic A. Network systems biology for drug discovery. *Clin Pharmacol Ther* 2010;**88**:120-125.
81. Alam-Faruque Y, Huntley RP, Khodiyar VK, Camon EB, Dimmer EC, Sawford T, *et al.* The impact of focused gene ontology curation of specific mammalian systems. *PLoS ONE* 2011;**6**:e27541
82. Khodiyar VK, Hill DP, Howe D, Berardini TZ, Tweedie S, Talmud PJ, *et al.* The representation of heart development in the gene ontology. *Dev Biol* 2011;**354**:9-17

## FIGURE LEGENDS

**Figure 1. Proteomics approaches.** Protein extracts can either be fractionated at the protein level prior to digestion or after protein digestion at the peptide level. In DIGE the protein extracts are labeled with different fluorescent dyes before they are separated by 2-DE. For SILAC, cells are metabolically labeled in culture by incorporation of heavy or light amino acids. Alternatively, labeling is performed at the peptide level, using iTRAQ or TMT isobaric tags. Peptides are then analysed by tandem mass spectrometry (MS/MS). 2-DE, two-dimensional gel electrophoresis; DIGE, difference gel electrophoresis; 1-DE, one-dimensional gel electrophoresis; SILAC, stable isotope labeling with amino acids in cell culture; AA, amino acid; iTRAQ, isobaric tag for relative and absolute quantitation; TMT, tandem mass tag.

**Figure 2. Gel-based proteomics.** Separation of the murine cardiac proteome by DIGE on different immobilized pH gradients: pH 3-10 NL **(A)** and pH 4-7 **(B)**. The white box highlights the better resolution of the same area on the narrow pH gradient. **(C)** Molecular weight distribution of 6 extracellular glycoproteins by SDS-PAGE. The characteristic “laddering” in abdominal aortic aneurysms (AAA) compared to normal aortic tissue (CON) is indicative of proteolysis. **(D)** Incubation of healthy aortic tissues with matrix metalloproteinases-12 (MMP-12) induced a similar fragmentation pattern of fibronectin as observed in AAA. In comparison, degradation by matrix metalloproteinases-9 (MMP-9) was less pronounced (reproduced with permission from Didangelos *et al*, Mol Cell Proteomics, 2011<sup>15</sup>).

**Figure 3. Computational approaches in proteomics.** Bioinformatics has become an essential part of the proteomic workflow to comprehensively analyse and visualize global changes in proteins as biological networks.

**Figure 4. Protein Identification.** The visualisation of extracellular matrix (ECM) proteins, and the corresponding correlation networks, identified by proteomics in the secretome of two murine cell types: **(A)** aortic smooth muscle cells (SMC) and **(B)** cardiac fibroblasts (CF). Correlation networks have been thresholded at a correlation coefficient  $> 0.90$ . **(C)** Number of ECM proteins identified. Number of nodes (proteins) **(D)** and links (correlations) **(E)** found in each cell type, as well as those in common.

**Figure 5. Metabolomics.** A comparison of control and hibernating murine hearts by high-resolution magic-angle-spinning  $^1\text{H}$ -magnetic resonance spectroscopy (HRMAS  $^1\text{H}$ -MRS) analysis from solid hearts **(A)** and  $^1\text{H}$ -nuclear magnetic resonance spectroscopy ( $^1\text{H}$ -NMR) of cardiac tissue extracts **(B)**. Both techniques showed consistent changes in metabolites, i.e. the ratio of glutamate, lactate, and taurine in hibernating compared to control hearts as determined by HRMAS  $^1\text{H}$ -MRS was 0.81, 1.09, and 0.79, respectively, which is in good agreement with the measurements of 0.68, 1.13 and 0.72 for the same metabolites by  $^1\text{H}$ -NMR. HRMAS  $^1\text{H}$ -MRS provides a means for measuring metabolites in intact hearts *ex vivo*.  $^1\text{H}$ -NMR of tissue extracts offers better resolution and allows the identification of more metabolites than HRMAS  $^1\text{H}$ -MRS spectra obtained from solid tissue.

Figure 1






Figure 2




Figure 3





Figure 4


A

Aortic Smooth Muscle Cells (SMC)




B

Cardiac Fibroblasts (CF)




C




Number of proteins identified

D





Number of nodes (proteins) in the correlation networks

E



Number of links (correlations > 0.9) in the correlation networks

Figure 5



**Table 1. Comparison of Proteomics Methods**

<b>Abbr.</b>	<b>Full name and explanation</b>	<b>Advantages</b>	<b>Disadvantages</b>
DIGE	Difference gel electrophoresis	Quantitation at the protein level Visualisation of posttranslational modifications and protein isoforms Good quantitative accuracy	Low sensitivity Only the differentially expressed proteins tend to be identified by MS/MS Proteins with very high or low pI or MW are not resolved on the gel
gel-LC-MS/MS	Separation by SDS-PAGE before LC-MS/MS analysis	Ease of use Prefractionation before LC-MS/MS analysis increases sensitivity "Laddering" as indication of proteolytic degradation	Prefractionation increases time requirements for MS/MS analysis Poor quantitative accuracy in complex mixtures without peptide labeling Proteins with very high or low MW are not resolved on the gel
SILAC	Stable isotope labelling with amino acids in cell culture	Minimal experimental variation Excellent quantitative accuracy Ease of use for cells in culture that proliferate and tolerate filtered serum supplements	Quantitation the peptide level Not suitable for cells that do not proliferate in culture, i.e. cardiomyocytes Metabolic labeling of animals is expensive
ITRAQ, TMT-tags	Isotopic labeling of peptides	Good quantitative accuracy Can be used with tissues as well as cell cultures	Quantitation the peptide level Mixed MS/MS spectra will contain reporter ions from different peptides

**Table 2. Differential Expression Methods**

<b>Abbr.</b>	<b>Name</b>	<b>Reference</b>
NSAF	Normalized Spectral Abundance Factor	Zybailov <i>et al.</i>
PLGEM	Power Law Global Error Method	Pavelka <i>et al.</i>
SI <sub>N</sub>	Normalized Spectral Index	Griffin <i>et al.</i>
SpI	Spectral Index	Fu <i>et al.</i>
Qspec	Qspec	Choi <i>et al.</i>
Hybrid	Hybrid-base approach	Wang <i>et al.</i>

**Table 3. Functional Annotation Databases**

<b>Abbr.</b>	<b>Name</b>	<b>Website</b>
BBID	Biological Biochemical Image Database	<a href="http://bbid.grc.nia.nih.gov/">http://bbid.grc.nia.nih.gov/</a>
BioCarta	BioCarta Pathways	<a href="http://www.biocarta.com/genes/index.asp">http://www.biocarta.com/genes/index.asp</a>
GO	Gene Ontology	<a href="http://www.geneontology.org/">http://www.geneontology.org/</a>
KEGG	Kyoto Encyclopedia of Genes and Genomes	<a href="http://www.genome.jp/kegg/">http://www.genome.jp/kegg/</a>
NCBI BioSystems	NCBI BioSystems Database	<a href="http://www.ncbi.nlm.nih.gov/biosystems">http://www.ncbi.nlm.nih.gov/biosystems</a>
OMIM	Online Mendelian Inheritance in Man	<a href="http://omim.org/">http://omim.org/</a>
PANTHER	Protein Analysis Through Evolutionary Relationships	<a href="http://www.pantherdb.org/">http://www.pantherdb.org/</a>
PID	NCI-Pathway Interaction Database	<a href="http://pid.nci.nih.gov/">http://pid.nci.nih.gov/</a>
Reactome	Reactome	<a href="http://www.reactome.org/">http://www.reactome.org/</a>
WikiPathways	WikiPathways	<a href="http://www.wikipathways.org/index.php/WikiPa">http://www.wikipathways.org/index.php/WikiPa</a>